

Ruh J, Ryschich E, **Schmidt J**, Secchi A, Gebhard MM, Herfarth C, Klar E  
Direkte Messung des Blutflusses an der Dünndarmschleimhaut der Ratte mit FITC-  
markierten Erythrocyten. Measurement of blood flow in the mucosa of the small intestine  
with FITC-labeled erythrocytes.  
Langenbecks Arch Chir 1997, 114 Suppl 1455-459

Leutloff UC, Eisold S, Schenk JP, Nöldge G, **Schmidt J**, Kauffmann GW  
[Obstruction in the duodenal passage. Mesenteric artery duodenal compression].  
Radialoge 2000, 40:83-85

Maksan SM, Ülger Z, Gebhard MM, Schmidt J 1.5  
Positiver Effekt von Antithrombin III auf die mesenteriale und hepatische  
Leukozytenadhärenz und Transaminasenanstieg bei entzündlicher  
Darmerkrankung und Leberzirrhose der Ratte. Positive effects of antithrombin  
III on mesenteric and hepatic leukocyte adherence and the course of liver  
enzymes in inflammatory bowel disease and liver cirrhosis in rats.  
Chir Forum 2002, 31:361-363

Leowardi C, Heuschen G, Kienle P, Heuschen U, Schmidt J 1.3  
Surgical treatment of severe inflammatory bowel diseases.  
Dig Dis 2003, 21:54-62

Autschbach F, Eisold S, Hinz U, Zinser S, Giese T, Linnebacher M, Löffler 6 9  
T, Büchler MW, Schmidt J  
High prevalence of *Mycobacterium avium* subspecies paratuberculosis IS900  
DNA in gut tissues from individuals with Crohn's disease.  
Gut 2005, 54:944-949

Maksan SM, Ulger Z, Gebhard MM, Schmidt J (2005) 3.3  
Impact of antithrombin III on hepatic and intestinal microcirculation in  
experimental liver cirrhosis and bowel inflammation: an in vivo analysis.  
World J Gastroenterol. 11:4997-5001

Maksan SM, Ryschich E, Ulger Z, Gebhard MM, Schmidt J (2005) 1 3.3  
Disturbance of hepatic and intestinal microcirculation in experimental liver  
cirrhosis.  
World J Gastroenterol 11:846-849

Mudter J, Weigmann B, Bartsch B, Kiesslich R, Strand D, Galle PR, Lehr HA, 6 5.6  
Schmidt J, Neurath MF (2005)  
Activation pattern of signal transducers and activators of transcription (STAT)  
factors in inflammatory bowel diseases.  
Am J Gastroenterol. 100:64-72

1.5

Holtmann MH, Krummenauer F, Claas C, Kremeyer K, Lorenz D, Rainer O, Vogel I, Bocker U, Bohm S, Buning C, Duchmann R, Gerken G, Herfarth H, Lügering N, Kruis W, Reinshagen M, Schmidt J, Stallmach A, Stein J, Sturm A, Galle PR, Hommes DW, D`Haens G, Rutgeerts P, Neurath MF

Long-term effectiveness of azathioprine in IBD beyond 4 years: a european multicenter study in 1176 Patients

Dig Dis Sci 2006, 51: 1516-1524

Antolovic D, Kienle P, Knaebel HP, Schmidt J, Gutt C, Weitz J,  
Koch M, Buchler MW, Seiler Ch

Totally laparoscopic versus conventional ileoanal pouch procedure--design of a single-centre, expertise based randomised controlled trial to compare the laparoscopic and conventional surgical approach in patients undergoing primary elective restorative proctocolectomy--LapConPouch-Trial.

BMC Surg 2006, 6:13

Welsch T, Hinz U, Loeffler Th, Muth, Herfarth Ch, Schmidt J,  
Kienle P

Early re-laparotomy for post-operative complications is a significant risk factor for recurrence after ileocaecal resection for Crohn's disease.

Int J Colorectal Dis 2007, 22:1043-9

Antolovic D, Koch M, Galindo L, Wolff S, Music E, Kienle P, Schemmer P, Friess H, Schmidt J, Buchler MW, Weitz J

Hepaticojejunostomy--analysis of risk factors for postoperative bile leaks and surgical complications.

J Gastrointest Surg 2007, 11:555-561