

## **Publikationen Dr. med. Thomas Scherer**

### **Originalarbeiten:**

Scherer Th., Kieser Ch., Gerber H.: Investigation and follow-up of 110 patients with monarthritis of unknown origin. Ther. Umsch. 46, 258 - 264, 1989.

Scherer Th., Furger F., Münch U.: Delay in hospital admission of patients with acute myocardial infarction. Schweiz. Rundschau Med. (Praxis), 78, 993 - 996, 1989.

Hines II M.E., Scherer T., Silwany O., Lauredo I.T., Wanner A., Stein-Streilein J., Altman N.H., Abraham W.M.: A glycolipid compound derived from *Mycobacterium avium* serovar 2 that inhibits the candidacidal activity of macrophages scavenges reactive oxygen species, but has no effect on nitric oxide production. Proceedings of the Fourth International Colloquium on Paratuberculosis 1995; 240-249, 1995.

Scherer T.A., Lauredo I.T., Abraham W.M. Scavenging of reactive oxygen species by a glycolipid fraction of *Mycobacterium avium* serovar 2. Free Radical Biolog. Med. 22:561-5, 1997.

Bingisser R., Kaplan V., Scherer T., Russi E.W., Bloch K.E. Effect of training on repeatability of cardiopulmonary exercise performance in normal men and women. Med. Sci. Sports Exerc. 29:1499-1504, 1997.

Scherer T.A., Barandun J., Martinez E., Wanner A., Rubin E.M. Effect of high frequency oral airway and chest wall oscillation and conventional chest physical therapy on expectoration in stable cystic fibrosis. Chest 113:1019-27, 1998.

Kurz D.J., Aguzzi A., Scherer T.A. Decompensated cor pulmonale as the first manifestation of adult-onset myopathy. Respiration 65: 317-319, 1998.

Fleisch M.C., Bläuer F., Gubler J.G.H, Kuhn M., Scherer Th.A.. Eosinophilic pneumonia and respiratory failure associated with Venlafaxine treatment. Eur. Respir. J. 2000; 15: 205-208.

Scherer Th. A. Nd-YAG laser ignition of silicone endobronchial stents. Chest 2000; 117: 1449-1454.

Scherer TA, Spengler C.M., Owassapian D., Imhof E., Boutellier U. Respiratory muscle endurance training in chronic obstructive pulmonary disease: impact on exercise capacity, dyspnea, and quality of life. Am. J. Respir. Crit. Care. Med. 2000; 162: 1709-1714.

Inderbitzi Rolf, Thomas Scherer, Jürg Barandun, Karl Klingler. NSCLC-Screening: Pro. Schweiz. Med. Forum 2003; 3: 656-60.

The International Early Lung Cancer Action Program Investigators. Computed tomographic screening for lung cancer. The relationship of disease stage to tumor size. Arch Intern Med 2006; 166: 321-5.

The International Early Lung Cancer Action Program Investigators. Women's susceptibility to tobacco carcinogens and survival after diagnosis of lung cancer. JAMA 2006; 296:180-4.

Henschke CI; for the International Early Lung Cancer Action Program Investigators. Survival of patients with clinical stage I lung cancer diagnosed by computed tomography screening for lung cancer. Clin Cancer Res. 2007; 13: 4949-50.

The International Early Lung Cancer Action Program Investigators. Survival of patients with stage I lung cancer detected on CT screening. N Engl J Med 2006; 355: 1763-71.

Abstracts:

Scherer Th., Münch U.: Etiology of dyspnea causing hospital admission. Schweiz. Med. Wschr. 119, Suppl. 28, 45, 1989.

Scherer Th., Lauredo I.T., Abraham W.M. Inhaled Interleukin 8 (IL-8) reduces neutrophil (PMN) and alveolar macrophage (AM) adherence and AM activation. Am. J. Respir. Crit. Care Med. 151 (suppl), A 137, 1995.

Güntert D., Imhof E., Scherer Th. Severe combined immunodeficiency syndrome due to incomplete (late onset) adenosine deaminase deficiency. Schweiz. Med. Wochenschr. 126 (suppl 75):10S, 1996.

Güntert DG., Mäder A., Scherer Th., Imhof E. Percutaneous embolotherapy of a pulmonary arteriovenous malformation with systemic arterial supply and venous drainage over pulmonary veins. Schw. Med. Wochenschr. 1997, 127, Suppl 84; 15S.

Scherer Th, Güntert D, Hofer M, Imhof E, Oelz O, Metzger U, Heitz M. Rigid vs. flexible bronchoscopy for endobronchial treatment. Am. J. Respir. Crit. Care. Med. 1999

Hofer M., Beynon Ch., Scherer Th., Göbel N., Brühlmann W., Lutz Th., Imhof E. Selektive Pulmonalisangiographie (SPA) versus Spiral-Angio-CT (SCT) bei Patienten mit klinischem Verdacht auf Lungenembolien (LE) und low oder intermediate Lungenperfusions- / Ventilationsszintigraphie (VPS). Schw. Med. Wochenschr. 1999

Scherer Th., Klingler K., Barandun J., Scheuer E., Went P., Metzger U., Inderbitzi R. Detektion von prämalignen Läsionen des Bronchialepithels: Weisslicht vs. Autofluoreszenz-Bronchoskopie. Forum Med. Suisse Suppl 1: 19S; 2001.

Summary and Discussions:

Scherer Th. Langzeiteffekte einer Prostazyklin-Dauerinfusion bei Patienten mit primärer pulmonaler Hypertonie. Schweiz Med Wochenschr 125:1408; 1995.

Metzger U., Scherer T. Das okkulte Bronchuskarzinom. In: Lungenkrebs, Fakten und Handlungsbedarf. Bundesamt für Gesundheit und Schweizerische Krebsliga. Nationales Krebs-Bekämpfungsprogramm, 70, 1997.