

Publikationsliste Dr. med. Hansjörg Vees

Vees H, Allal AS. Carbogen breathing combined with radical radiotherapy in advanced head and neck cancer patients with severe co-morbidities. *Clin Oncol (R Coll Radiol)*. 2006 Aug;18(6):493-6.

Miralbell R, **Vees H**, Lozano J, Khan H, Hidalgo A, Mollà M, Linero D, Rouzaud M. Endorectal MRI assessment of local relapse after surgery for prostate cancer: a model to define treatment field guidelines for adjuvant post-prostatectomy radiotherapy in patients at high-risk for local failure. *Int J Radiat Oncol Biol Phys*. 2007 Feb 1;67(2):356-61.

Vees H, Buchegger F, Albrecht S, Khan H, Hidalgo A, Mollà M, Husarik D, Hany TF, Miralbell R. ¹⁸F-choline and ¹¹C-acetate positron emission tomography (PET) in patients with detectable or increasing PSA after radical prostatectomy: a comparative assessment. *BJU Int*. 2007 Jun;99(6):1415-20.

Vees H, Mach N, Hügli A, Balmer Majno S. Is there a place for low-dose chest and prophylactic brain irradiation in limited-disease small cell lung cancer, *Med Oncol*. 2009;26(3):298-302.

Vees H, Senthamizhchelvan S, Ratib O, Weber DC, Miralbell R, and Zaidi H. Assessment of various strategies for delineation of target volumes in ¹⁸F-FET PET studies of high-grade gliomas patients. *Eur J Nucl Med Mol Imaging*. 2009 Feb;36(2):182-93.

Zaidi H, **Vees H**, Wissmeyer M, Molecular PET/CT imaging-guided radiation therapy treatment planning. *Acad Radiol*. 2009 Sep;16(9):1108-33.

Steiner, C, **Vees, H.**, Zaidi H., Berrebi, O., Kossovsky M.P., Khan G.H., Miralbell R., Ratib O., Buchegger F. Three-phase ¹⁸F-fluorocholine PET/CT in the evaluation of prostate cancer recurrence. *Nuklearmedizin*. 2009;48(1):1-9.

Wang H, **Vees H**, Miralbell R, Wissmeyer M, Steiner C, Ratib O, Senthamizhchelvan S, Zaidi H. ¹⁸F-fluorocholine PET-guided target volume delineation techniques for partial prostate reirradiation in local recurrent prostate cancer. *Radiother Oncol*. 2009 Nov;93(2):220-5.

Vees H, Bigler R, Bieri S, Gruber G. Well-known and new parameters in breast radiotherapy treatment planning: their correlation with irradiated heart volume. *Cancer Radiother*. 2011 Jul 25.

Schick U, Bolukbasi Y, Abdah-Bortnyak R, Kuten A, Thariat J, Caglar H, Ozsaran Z, Schleicher U, Lössl K, Zwahlen D, Villette S, **Vees H**. Outcome and prognostic factors in endometrial stromal tumors: A rare cancer network study. *Int J Radiat Oncol Biol Phys*. 2012 Apr 1;82(5):e757-63.

Vees H, Steiner Ch, Dipasquale G, Chouiter A, Zilli T, Velazquez M, Namy S, Ratib O, Buchegger F, Miralbell R. Target volume definition in high-risk prostate cancer patients using sentinel node SPECT/CT and ¹⁸F-choline PET/CT. *Radiat Oncol*. 2012 Aug 8;7(1):134.

Vees H, Casanova N, Zilli T, Imperiano H, Ratib O, Popowski Y, Wang H, Zaidi H, Miralbell R. Impact of ¹⁸F-FDG PET/CT on target volume delineation in recurrent or residual gynaecologic carcinoma. *Radiat Oncol*. 2012 Oct 22;7(1):176.

Vees H, Dipasquale G, Nouet P, Zilli T, Cozzi L, Miralbell R. Pelvic Lymph Node Irradiation including Pararectal Sentinel Nodes for Prostate Cancer Patients: Treatment Optimization comparing Intensity Modulated X-rays, Volumetric Modulated Arc Therapy, and Intensity Modulated Proton Therapy. *Technol Cancer Res Treat* 2014.