

open studies

	<b>name*</b>	<b>description</b>
<b>Breast cancer</b>	IMPASSION 030-Studie (BIG 16-05) POLAR-Studie (IBCSG 59/19) SAKK 21/18 Studie TARGIT B -Studie (UCL London) TAXIS-Studie (SAKK 23/16) TOUCH-Studie (IBCSG 55-17) WISE-Studie (IBCSG 95-17)	Atezolizumab (anti-PD-L1-AK) in combination with adjuvant chemotherapy (anthracycline/taxane-based) v.s. adjuvant chemotherapy alone for operable triple-negative breast cancer Palbociclib in HR-positive/HER2-negative, isolated, locoregional recurrence of breast cancer Ribociclib-endocrine combination therapy versus chemotherapy as first-line treatment in patients with visceral metastatic breast cancer, max. 55 patients in CH over 2y Radiotherapy study: Intraoperative boost v.s. conventional boost Targeted versus radical axillary lymph node surgery followed by radiotherapy in patients with axillary lymph node metastasis (es) Neoadjuvant palbociclib in combination with letrozole and dual Her2 blockade v.s. Paclitaxel in combination with dual Her2 blockade for older patients > 65 years with hormone-sensitive, Her2-negative early breast cancer. 24-week activity programme for patients with early breast cancer undergoing aromatase inhibitor therapy
<b>Urogenital tumours</b>	PCO Studie (Prostate Cancer Outcome) SAKK 08/15 Studie KSSG SAG TCCS Studie SAKK 06/17 Studie	Measuring the quality of life of prostate cancer patients using a globally standardised patient questionnaire The study examines the efficacy and tolerability of the drug Metformin in combination with radiation in prostate cancer patients with recurrence after prostate removal Swiss Austrian German Testicular Cancer Cohort Study/Observational Study, the aim of which is to improve the quality of care for patients with testicular cancer in Switzerland. multicentre, single-arm phase II study: Immunotherapy in combination with standard neoadjuvant chemotherapy in patients with operable urothelial carcinoma
<b>Gastrointestinal tumours</b>	Roche GLYCART Studie SAKK 41/14 - ACTIVE-2 Studie SAKK Prospect-Trial (N1048) PIPAC/PITAC Studie  Dante Studie (Flot8) PREFER-Studie Barret-CAD Studie	Ex vivo model of fresh tumour tissue Effect of physical activity in patients with metastatic colorectal Ca A phase II/III study comparing pretreatment of locally advanced rectal cancer with either combined radio-chemotherapy or chemotherapy alone, always followed by surgical tumour removal as total mesorectal excision (TME). Efficacy and safety of pressurized intraperitoneal / intrathoracal aerosol chemotherapy (PIPAC / PITAC) with cisplatin and doxorubicin in gastric, ovarian and colorectal cancer with peritoneal carcinomatosis and oxaliplatin in patients with mesothelioma / pleural cancer with pleural carcinomatosis. A prospective single center observational study. A randomized, open-label Phase II efficacy and safety study of atezolizumab in combination with FLOT versus FLOT alone in patients with gastric cancer and adenocarcinoma of the oesophago-gastric junction (MO30039) Endoscopic management of patients with T1bN0M0 esophageal adenocarcinoma Data collection for computer aided detection of Barrett neoplasia
<b>Hematological tumours</b>	EBMT-Register SAKK 35/14 Studie SAKK 36/13 Studie SAKK 39/16 OptiPOM Studie SAKK HD21 Studie Triangle Studie	Register for patients undergoing haematopoietic stem cell transplantation or other cell therapy (e.g. CAR-T cell therapy) Rituximab with or without Ibrutinib for untreated patients with advanced follicular lymphoma in need of therapy. A randomized, double-blinded, SAKK and NLG collaborative Phase II trial Combination of ibrutinib and bortezomib followed by ibrutinib maintenance to treat patients with relapsed and refractory mantle cell lymphoma. A multicenter Phase I/II trial. investigates ways to improve the treatment of advanced multiple myeloma Treatment optimization trial in the first-line treatment of advanced stage Hodgkin lymphoma; comparison of 6 cycles of escalated BEACOPP with 6 cycles of BrECADD. Autologous stem cell transplantation (ASCT) after Rituximab/Ibrutinib/Ara-C treatment in patients with mantle cell lymphoma
<b>Gynaecology tumours</b>	PIPAC/PITAC Studie  SENTIX (SENTinel lymph node in cervIX cancer) Studie Roche GLYCART Studie	Efficacy and safety of pressurized intraperitoneal / intrathoracal aerosol chemotherapy (PIPAC / PITAC) with cisplatin and doxorubicin in gastric, ovarian and colorectal cancer with peritoneal carcinomatosis and oxaliplatin in patients with mesothelioma / pleural cancer with pleural carcinomatosis. A prospective single center observational study. a prospective observational trial on sentinel lymph node biopsy in patients with early stage cervical cancer Ex vivo model of fresh tumour tissue
<b>Observational studies (cros</b>	Consilium-Studie Symptomerfassung mittels APP  Alpine TIR/SAKK 80/19	Observational study: Symptom recording using an app. The study examines the use of a smartphone app with which patients can record changes in their well-being and the symptoms that occur during therapy. The primary aim of this is to examine the effects of different periods of time between the planned doctor's visits on the quality of the symptom recording. Swiss Tumor Immunology Registry (SwissTIR): Quality register for immuno-oncology
<b>Palliative Care</b>	SAKK 96/12 Studie (Reduse-Studie)	Palliative care/prostate or breast cancer – metastatic, castration-resistant (prostate) and with at least 3 bone metastases/therapy of patients with bone metastases with Xgeva®

\*study names are mentioned in their original language