contact:

cancer centre Hirslanden Zürich Witellikerstrasse 40 8032 Zürich Phone +41 44 387 96 62 tumorzentrum.hirslanden@hirslanden.ch

open studies

breast cancer name of study
CAMBRIA-2 study

EKIB study

EXPERT- study (ANZ 1601/ BIG 16-02)

MELODY-study PROMs

TAXIS-study (SAKK 23/16) TroFuse-010 study

VISION-study (SAKK 23/18)

urogenital tumors PCO study (Prostate Cancer Outcome)

MRI bladdercancer study SAKK 08/23 study

gastrointestinal tumors BERING CRC study

EDIUM study

DANTE Study

ESODATA study

GUIDE.MRD-01-MRC study PREFER-study

Quality assurance study for duodenal tumors

Quality assurance study for pancreatic tumors

Quality assurance study for

Tiger study

hematological systemic diseases Abbvie Valor-Studie

EBMT-Register PASS study

SAKK 38/23 Liberty study SAKK EMCL-Registry Treg-cell study

neuro-oncological tumors LGG Studie

Sinonasale QoL Studie

multi-organ studies AGE-1 Studie

Roche GLYCART Studie

target group/ short description

An adjuvant endocrine-based therapy study of camizestrant (AZD9833) in ER+/HER2-early breast cancer (CAMBRIA-2).

Analysis of human breast tumor heterogeneity and immune cell infiltrates and their relevance for patient outcome by mass cytometry.

Cinical trial which presents a unique opportunity to improve personalised use of radiation therapy in patients with early breast cancer according to their individual risks of local recurrence (molecular luminal A-breast cancer).

A prospective, non-interventional, multicenter cohort study to investigate various imaging-guided techniques for the localization of malignant breast lesions.

Patient reported outcomes after oncoplastic breast conserving surgery.

Tailored AXIIIary Surgery with or without axillary lymph node dissection followed by radiotherapy in patients with clinically node-positive breast cancer (TAXIS).

A study of Sacituzumab Tirumotecan as a single agent and in combination with Pembrolizumab versus treatment of physician's choice in participants with HR+/HER2- unresectable locally advanced or metastatic breast

Vacuum assisted biopsy immediately before surgery as an intra- or pre-operative surrogate for patient response to neoadjuvant chemotherapy for breast cancer (VISION I).

Patient reported outcome measrement of patients with prostate cancer with global uniform patient questionaire.

Multiparametric Magnetic Resonance Imaging (MRI) to improve bladder cancer staging and treatment decision-making.

Addition of Darolutamide to first line treatment of mCRPC: a randomized open label phase II trial".

Encorafenib and Cetuximab in patients with metastasized BRAFV600E mutated colorectal cancer: a multi-centric, multi-national, prospective, longitudinal, non-interventional study.

Patient-reported outcome study for colon cancer patients. Ergebnisqualität bei Darmkrebs: Identifikation von Unterschieden und Massnahmen zur flächendeckenden Qualitätsentwicklung (Quality of Experience in the case of colon cancer: identification of differences und actions to be taken for a comprehnesive quality development)

A randomized, open-label phase II/III study on the efficacy and safety of Atezolizumab in combination with FLOT versus FLOT alone in patients with gastric cancer and adenocarcinoma of the gastroesophageal junction with high immune reactivity.

International standardized online database für outcomes of the patients receiving evaluation and/ or treatment für esoohagal diseases of benign and malignant conditions.

Evaluation of ctDNA for the detection of minimal residual disease (MRD) after curative resection of colorectal cancer (CRC).

Endoscopic management of patients with T1bN0M0 esophageal adenocarcinoma.

Quality assurance study of perioperative morbidity and post-surgical follow up of hepato-duodeno-pancreatic surgery.

Quality assurance study of perioperative morbidity and post-surgical follow up of pancreatic surgery.

Robot-assisted minimally invasive thoraco-laparoscopic esophagectomy versus minimally invasive esophagectomy for resectable esophageal adenocarcinoma, a randomized controlled trial.

Distribution of lymph node metastases in esophageal carcinoma.

A study to assess change in disease state in adult participants with acute myeloid leukemia (AML) ineligible for intensive chemotherapy in Switzerland and Austria receiving oral Venetoclax tablets (VALOR).

Registry for patients that received a haematopoietic stem cell transplantation or CAR T-cell therapy.

Observational study including patients treated with CAR-T therapy to assess treatment outcomes and associated complications.

Liquid Biopsy to diagnose and monitor CNS involvement in High-risk B Cell Non-Hodgkin Lymphoma

The registry of the European Mantle Cell Lymphoma study group - MCL: Disease pattern, treatment algorithms and mechanisms of relapse.

Regulatory T-cells as tumor-promoting cellular partners and potential therapeutic targets in diffuse large B-cell lymphoma.

Prospective observational study on guideline-based therapy in astrocytomas and oligodendrogliomas, WHO grade 2, with IDH1/2 mutation.

Prospective observational study to evaluate the nasal quality of life after endoscopic transnasal surgeries.

Evaluation of forensic methods for age estimation of patient with cancer and a cotrol group.

Ex vivo modeling of fresh tumor tissue